

RESEARCH

Open Access



An ethnobotany survey of wild plants used by the Tibetan people of the Yadong River Valley, Tibet, China

Chang-An Guo^{1,2†}, Xiao-Yong Ding^{1,2†}, Yi-Won Addi^{1,2}, Yu Zhang¹, Xiao-Qian Zhang^{1,2}, Hui-Fu Zhuang¹ and Yu-Hua Wang^{1*}

Abstract

Introduction: Plant resources gathered from the wild are important sources of livelihood needs, especially for low-income populations living in remote areas, who rely on these plants for food, fuelwood, medicine and building materials. Yadong County is a valley at the border between the China, India and Bhutan in southern Tibet. Yadong is rich in biodiversity and culture, but ethnobotanical knowledge has not been systematically studied. This study aimed to document the ethnobotanical knowledge of Tibetans in Yadong County.

Methods: Ethnobotanical data were documented through free listings, key informant interviews, and semi-structured interviews during fieldwork. The culture importance index (CI) and informant consensus factor index (FIC) were used as the quantitative indices.

Results: In total, 163 informants (46 women and 117 men). A total of 3,031 use reports and 121 plant species belonging to 52 families and 91 genera were included. These use reports were then classified into 20 categories belonging to 9 major categories. The utilisation category that containing the most plant species was food, followed by economic, medicine, animal feed, social uses, other uses, environmental uses, materials and fuels. Among the economic plants, 32 medicinal plants are traditionally used in the local region. Plants with high CI included *Fritillaria cirrhosa*, *Neopicrorhiza scrophulariiflora*, *Betula utilis*, *Rheum nobile*, and *Urtica hyperboreana*.



Conclusion: This research demonstrates the diversity of the types and functions of Yadong Tibetan traditional plant knowledge. Knowledge of edible and medicinal plants in this area is prominent, reflecting the ability to cope with the lack of fruits and vegetables and basic family medical care. There were exchanges between the traditional plant culture in the study area and its surroundings. With socioeconomic development, the commercial value of medicinal plants has increased, and locals are also seeking ways to adopt sustainable development to cope with the excessive consumption of plant resources.

Keywords: Eastern Himalayas, Biodiversity hotspots, Tibetan, Traditional knowledge, Yadong county, Culture exchange

*Correspondence: wangyuhua@mail.kib.ac.cn

†Chang-An Guo and Xiao-Yong Ding contributed equally to this work

¹ Yunnan Key Laboratory for Wild Plant Resources, Kunming Institute of Botany, Chinese Academy of Sciences, 132# Lanhei Road, Yunnan 650201, Kunming, China

Full list of author information is available at the end of the article

Introduction

Wild plant resources are important sources of food, fuelwood, medicine, forage and building materials, especially for the poor, living in remote areas [1]. However, the traditional understanding of wild plants is rapidly lost due to the development of socio-economic development [2].



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Moreover, traditional knowledge depends on specific locations and information passed down through generations [3]. Thus, documentation and evaluation of traditional ethnobotanical knowledge are urgently needed [4].

Yadong County is located on the edge of biodiversity hotspots in the eastern Himalayas, India, and Myanmar [5]. The Yadong River runs through Yadong County and presents a valley topography; Yadong County is also called Yadong River Valley and is an important border area in southern Tibet bordering Bhutan and Sikkim to the east, west, and south and has 41 mountain passes connecting Yadong to Bhutan and Sikkim. Because of its proximity to the sea and its superior border trade environment, it has become the largest port for border trade in Tibet. In addition, trade between Nepal, India and China is active [6].

Tibetans live mainly on the Qinghai-Tibet Plateau. They have a long history and rich traditional cultural knowledge in many aspects such as food, medicine, religion, architecture, and handicrafts [7]. Although Tibetans are classified in China as a minority, they consist of several Tibetan languages, dialects and ways of life, with six main groups/dialects, three of which occur in China and are classified as Ü-Tsang, Kham and Amdo [8]. The language used by Tibetans in Yadong belongs to the Ü-Tsang dialect in the Tibetan branch of the Tibeto-Burman language group [9]. The main sources of income for the local population are livestock rearing and gathering herbs. Nomads seasonally migrate to high altitudes during the summer with their yaks and return to permanent settlements before the onset of the winter. Herbal gatherers usually harvest during spring and summer when plants grow. Seasonal labour across altitudes is part of their lives.

The main Tibetan settlements abroad are in India, Nepal, and Bhutan, most of which are in the southern foothills of the Himalayas. Foreign ethnobotanical research is mainly concentrated in these countries and regions, and related research involves the utilisation of edible plants, medicinal plants, handicrafts, dyeing, feed, and fibres. Tibetans live in vast area of China, including Tibet, Qinghai, western Sichuan, and northwestern Yunnan. However, ethnobotany research on Tibetans in China is mainly concentrated in some provinces and cities in the eastern Tibetan region. In most Tibetan areas, ethnobotanical research is still lacking.

Tibetans have a rich and unique knowledge of the local environment, such as climate, soil, wildlife, vegetation, and plant utilisation [10]. This traditional knowledge stems from the interrelationships among humans, plants, animals, natural phenomena, and religious beliefs. Predecessors have conducted surveys in Tibet and other areas where Tibetans live and have found that some wild plants

have provided many products and services for locals. Traditional Tibetan medicine has been the main disease treatment in many remote areas, relying on a large number of wild plants as well as traditional treatment methods [11–30], and some of these medicinal plants are also collected for trade [26, 28, 31–34]. Various wild edible plants have been collected for food [23, 24, 32, 33, 35–41]. Tibetans also place flowers collected from the wild on the altars of houses and temples or collect Tibetan incense plants to worship their gods [25, 42]. In addition, there is a wealth of knowledge on forage, house building, tools, fuels, dyes, and seasoning [11, 25, 43].

However, to date, traditional knowledge of plants used by Tibetans in Yadong has not been explored. Therefore, this research aimed to document the ethnobotanical knowledge of Tibetans in Yadong County.

Method


Study area

Yadong County, which belongs to Shigatse City, Tibet, is located in the southern Himalayan mountains and borders India and Bhutan (Fig. 1). The altitude ranges from 1,600 to 7,300 m, and the annual average temperature is 0 °C, with the average temperatures of the coldest and hottest months of -5 °C and 10°C, respectively [44]. The terrain is high in the north and low in the south. The northern part of Yadong is an important plateau pasture in Tibet, and is mainly composed of alpine ecosystems. The southern part of Yadong has a large area of virgin forest, which has a mild climate and abundant water resources, and is considered a green treasure house in the Himalayas. The total population of Yadong County is 13,992, among which Tibetans account for 98%, Han 1.5% and other minorities 0.5% [44]. Economically, the Yadong District can be defined as a rural area based on agricultural and livestock activities. *Hordeum vulgare* var. *coeleste* is the predominant crop, and *Bos grunniens* is the main livestock [45–47].

Data collection

In August 2020 and May 2021, our ethnobotany fieldwork was conducted on 12 villages of 7 townships in Yadong (Fig. 1). First, field study permission was obtained from the local community committee and government authority. We explained our purpose to local governments and requested assistance from them. All our fieldwork was conducted with informed consent.

The snowball sampling method was used to select the key informants, such as veterinarians and herdsmen. Other informants were selected by the randomized household interview method. In total, traditional knowledge was collected from 163 informants. Ethnobotanical knowledge was collected by semistructured



face to face interviews. Because many Tibetans in the study area cannot speak Mandarin fluently, the field work was performed with the assistance of local guides who were employed with the help of local community leaders. All interviews were conducted in the Tibetan language, which was translated into Mandarin by local guides. All field studies were conducted with the consent of informants. According to the commonly used 5 w + 1 h (What, Where, When, Who, Why, How) principle in ethnobotany, this study designed the following questions for semistructured interviews:

1. Would you mind listing some wild plants you have used?
2. How to use this plant?
3. Which plant parts were used, roots, stems, leaves or other parts?
4. Why do you use this species?
5. What time do you collect this plant?

The questions were designed to collect data on the (i) vernacular name of the plants, (ii) category of use, (iii) parts used, (iv) methods for preparation and administration, (v) characteristics of the plant material (dried or fresh) and (vi) collection time.

The specimens were collected from the field of survey with the help of the key informants and all materials are labelled with numbers and names. Photographs of each plant were taken. All specimens were kept in the herbarium of Kunming Institute of Botany (KUN). The Flora of China was used to help identify the plants [48] and The Plant List was used to ensure the Latin name of the plants [49].

Data analysis

We adopted the Use report (UR), cultural important index (CI) and informant consensus factor index (FIC) as ethnobotanical indices. All information about the use of local plants was organized into a “use report”

list consisting of three parts: informant, used plant and used category [50, 51].

The cultural important index (CI) [52] was the sum of the proportion of informants that mentioned each of the use categories for a given species. This index is used to quantitatively evaluate the importance of a certain plant to Yadong Tibetans from the perspective of comprehensive value. In other words, CI represents the diversity of plant uses and the degree of recognition of information sources for each use category. The calculation formula is as follows:

$$CI = \sum_{U=u1}^{uNC} \sum_{i=i1}^{iN} \frac{URui}{N}$$

NC was the total number of use categories and N was the total number of informants. CI ranges between 0 and the number of all utilization categories. A higher CI value indicated the multiple uses of a species and a higher degree of recognition.

The informant consensus factor index (FIC) was developed by Robert T. Trotter [53]. FIC was used to evaluate the degree of consensus among the population about how to treat a particular disease. The calculation formula is as follows:

$$FIC = \frac{Nur - Nt}{Nur - 1}$$

where Nur is the number of use reports from the informants for a particular disease and Nt is the total number of plant species used to treat the disease. The FIC values range between 0 and 1. A higher FIC means that different herbalists have a higher consensus on the plant species used to treat certain diseases.

Results

Distribution of knowledge among informants

This study documented 3,028 use reports (UR) from 163 informants. Among the informants, 46 were women and 117 were men (Table 1). The selection of our information reporter was random but resulted in more men and fewer women. This reason may be because the right to speak is mainly in the hands of men, who are primarily responsible for external affairs and livelihoods in most families, while women are mainly responsible for household affairs [54]. The informants were aged from 7 to 81 years, the average age was 52 years, while that of men was 53 years, and 49 years for women. Middle-aged individuals (40–59 years) provided the most use reports on plant use, whereas young individuals under 30 years old and the elderly over 70 years old provided fewer Urs than did middle-aged people (Fig. 2).

Taxonomic diversity of wild plants used by locals

In total 121 plant species belonging to 52 families and 91 genera were identified in the study area. The most cited family was Compositae (16 species), followed by Rosaceae (9), Polygonaceae (7), Ericaceae (6), Apiaceae (5) and Lamiaceae (5) (Table 2). Among the plants, 89 were herbaceous, 20 were shrubs, 10 were trees, and 2 were vines (Table 2).

In our survey, the most frequently used parts were whole plants (37), followed by fruits (27), roots (27), leaves (23), stems (16), aerial parts (13), flowers (12), bulbs (1), seeds (1), and burls (1) (Table 2).

The diversity of use categories

The informants referred to 3,028 URs, which 1,177 (38.8%) were for food uses, 805 (26.6%) for medicinal uses, 560 (18.5%) for economic plants and 486 (16.1%) for other categories. There were 53 edible plants species, 53 economic plants, 46 medicinal plants, and 50 plants used in other categories, including animal feed (18), social uses (12), environmental uses (6), materials (5), fuel (4), and other uses (10) (Table 3). Many plants belonged to multiple utilisation categories. There were 30 plant types plants with three or more uses: 12 medicinal and edible homologous plants, 32 plants that could be used both as a source of income and as medicinal plants, and 15 plants that could be used as both a source of income and food (Table 2).

Food

A total of 53 species of wild plants are used as food for humans. The food categories in Yadong include fruits (28 species), vegetables (16), seasoning (7), starches (3), and beverages (1) (Table 3). The most widely used component is the fruit. There are very few gardens and woodlands available for planting fruit trees in the local area, and the yield of local fruit tree varieties is extremely low. This may be one of the main reasons why the locals collect wild fruits from the wild as a nutritional supplement.

Animal feed

Eighteen wild plant species were used as animal feed. Among these 18 species, 16 were herbs, and 2 were woody plants (Table 3). These included *Heracleum nyalamense* ($CI_{animal\ food}=0.1840$), *Thermopsis barbata* (0.0798), *Polygonum macrophyllum* (0.0613), *P. tortuosum* (0.0552), and *Cirsium eriophoroides* (0.0429). *H. Nyalamense* is the most popular animal feed plant for locals. Local people indicated that cattle grew more strongly after feeding on this grass.

Table 1 Characteristics of informants

Characteristics	Number	Percentage
<i>Communities</i>		
Qudancha	13	8.0
Guru	12	7.4
Duina	13	8.0
Pari	17	10.4
Shangkangbu	18	11.0
Xiakangbu	17	10.4
Gangu	17	10.4
Rubingang	10	6.1
Galingang	9	5.5
Zhuju	8	4.9
Chema	10	6.1
Renqingang	19	11.7
<i>Gender</i>		
Female	46	28.2
Male	117	71.8
<i>Age</i>		
Below 29	15	9.2
30–39	26	16.0
40–49	22	13.5
50–59	42	25.8
60–69	39	23.9
Above 70	19	11.7

Medicinal plant use

A total of 43 traditional medicinal plants belonging to 24 families and 39 genera have been documented to treat 14 different types of human diseases, including dermatologic disorders, gastrointestinal problems, respiratory diseases, diarrhoea, and arthritis (Table 4). The most cited families of medicinal plants were Compositae (7 species), followed by Polygonaceae (4), Gentianaceae (4), and Lamiaceae (3). The plant parts most commonly used for remedy preparation are the roots accounting for 45.5% of the total medicinal plants. In addition, 10 veterinary medicines were used to treat 5 types of animal diseases (Table 4). The five most-cited species were *Gentiana veitchiorum* ($CI_{medicine} = 0.5767$), *Neopicrorhiza scrophulariiflora* (0.5215), *Fritillaria cirrhosa* (0.4969), *Taraxacum tibetanum* (0.3436), and *Fraxinus paxiana* (0.3006) (Table 2).

The FIC of 14 diseases ranged from 0.67 to 1, and were the highest for toothaches (1.00) and hypoimmunity (1.00), followed by inflammation (0.98), skeleto-muscular system disorders (0.98), and respiratory complaints (0.97). The FIC values were lowest for infections (0.67). The most cited diseases were respiratory complaints (321 URs), followed by inflammation (142), gastrointestinal (118) and dermatopathy (116) (Table 4).

Economic plants

Yadong County is rich in medicinal and non-timber forest products. We found 53 commercial plants in the study area. These plants were mainly sold to Tibetan doctors and Hui merchants. Among them, 32 species were used as medicinal plants by locals (Table 3). The most frequently mentioned economically important plant was *Fritillaria cirrhosa* ($CI_{economic} = 0.3374$), followed by *Saussurea tridactyla* (0.3313), *Rhodiola himalensis* (0.3252), *Angelica paeonifolia* (0.2454), and *Panax pseudoginseng* (0.1779) (Table 2). In addition, local people also collect *Cordyceps sinensis* or *Exidia* sp. as important sources of income.

In the local region, except for a small number of wild economic plants that collectors directly use, most plants enter the local or foreign market in somehow (Fig. 3).

Social uses

Social uses were divided into two categories: ritual plants (11 species), and tobacco substitutes (1) (Table 3). Tibetan people convey their wishes to the gods through various sacrificial activities and offer them several items, praying for happiness and well-being. A total of 11 species of plants are used in social activities. Among them, nine species were used for incense. For example,




Fig. 2 Distribution of knowledge among informants: The line represents the average number of URs provided by local people in each age group

Rhododendron anthopogon ($CI_{Social\ uses} = 0.4417$), *Nardostachys jatamansi* (0.2147), and *Juniperus indica* (0.1840). Two species were used at funerals. The local people use the purple pigment from *Onosma hookeri* (0.0675) roots to decorate the offerings. After a person dies, *Myricaria rosea* (0.0123) sticks are burned to pay homage to them.

Environmental uses

In the local region, milk collection showed seasonal characteristics and wild plants can be used for climate prediction [55]. Three plant species were mentioned in the interviews; *Primula sikkimensis* was the most frequently mentioned, followed by *P. concinna* and *Caltha palustris*. The yield and quality of yak milk are high when *P. sikkimensis* and *C. palustris* are in bloom. The flowering of *P. concinna* heralds the arrival of the rainy season and is also the time for planting and grazing. These plants have some common characteristics. These plants tending to grow in around pastures and farmlands. In addition, flowering was considered a climate predictor by most respondents.

Materials and other uses

Five plant types are used as raw materials for dyeing (3), papermaking (1), and crafts (1) (Table 3). However, the frequency of mentioning this type of utilisation was very low, which may indicate that the local area is losing traditional handicraft knowledge or that this knowledge is in the hands of only a few people. *Rheum acuminatum*, *Rheum nobile*, and *Polygonum tortuosum* are used in traditional dyeing. *R. acuminatum* and *R. nobile* roots are crushed, boiled in water, and used as yellow dyes. *Stellera*

chamaejasme is used to make paper, which is insect repellent, antiseptic, and flexible. The old stems of *Aristolochia griffithii* were picked up by locals and polished into ornaments or stools.

A total of 10 kinds of plants were used for other purposes (9 tools, and 1 repellent). Five plant species were used as make cooking tools (Table 3). For example, *Betula utilis* sticks can be used to make spoons or shovels. *Potentilla fruticosa* var. *arbuscula* sticks are used to make brooms for washing pots. *Enkianthus deflexus* branches are used to make a blender, which is used to prepare milk tea. Local Tibetans prefer to eat a fermented milk product (*pilu*), which has a unique taste, and whose fermentation process is notable. The locals collect *Betula utilis*, *Salix myrtillacea*, and *Salix daltoniiana* branches, boil, peel, and place them in a bucket, after which raw milk is poured over them and sealed to prevent mosquitoes from entering. This bucket is rotated and shaken daily so that raw milk is evenly attached to the branches. After 15 days, the local speciality food *pilu* (smelly cheese) is produced. *Pilu* must be cooked with yak butter before it can be eaten (Fig. 4). In the process of producing *pilu*, there is a critical step: placing branches in the bucket. If branches are not added, water loss will be slow and the milk will spoil. When asked why they chose the three trees mentioned above, the locals gave the following answers: they were easier to obtain, nontoxic and easier to peel their bark; most importantly, their branches were not easily corroded and could be reused many times.

Despite of social changes, some crafts and tools are still handmade and use plant materials [56]. These

Table 2 The wild plants used by the Yadong Tibetan

Botanical family	Local Tibetan script name (Latin form)	Botanical taxon	Voucher number	Local use (parts used) (UR)	CI
Adoxaceae	ན་ଲିମୋ (na-li-mo)	<i>Viburnum grandiflorum</i> Wall. ex DC.	QTP-EBT-4013	Food (fruits): raw, fresh fruit (16)	0.0982
Amaranthaceae	ଲେହ (Leh)	<i>Chenopodium album</i> L.	QTP-EBT-3073	Food (leaves): cooked vegetable (40)	0.2454
Amaryllidaceae	ରିଶ୍କୋ (ri-skho)	<i>Allium fasciculatum</i> Rendle	QTP-EBT-3050	Food (whole plants): cooked vegetable (63)	0.3926
Amaryllidaceae	ରିଶ୍କୋ (ri-skho)	<i>Allium prattii</i> C.H.Wright	QTP-EBT-3009	Food (whole plants): cooked vegetable (32)	0.1902
Amaryllidaceae	ତେଣ୍ବୁ (tsen-bu)	<i>Allium przewalskianum</i> Regel	QTP-EBT-3200	Food (leaves): seasoning (31)	0.1902
Amaryllidaceae	ଶକ୍ହୋଲାମୋ (skho-skho-lha-mo)	<i>Allium wallichii</i> Kunth	QTP-EBT-3108	Food (whole plants): cooked vegetable (29)	0.1779
Apiaceae	ତଙ୍କଖୋ (tang-khoe)	<i>Angelica paeoniifolia</i> R.H.Shan & C.C.Yuan	QTP-EBT-3077	1. Medicine (roots): soak in wine, stew in soup, to nourish the body (11) 2. Economic (roots) (40)	0.3006
Apiaceae	ଗୋମନ୍ଯେ (go-mnye)	<i>Carum carvi</i> L.	QTP-EBT-3137	1. Food (leaves and fruits): cooked vegetable; seasoning (40) 2. Economic (leaves and fruits) (2)	0.2577
Apiaceae	ତଙ୍ଗ (tong)	<i>Heracleum nyalamense</i> Shan & T.S. Wang	QTP-EBT-3075	Animal food (whole plants) (30): fodder	0.1840
Apiaceae	ଜା (Ja)	<i>Trachydium subnudum</i> C.B. Clarke ex H. Wolff	QTP-EBT-3206	Food (tender leaves): seasoning (21)	0.1288
Apiaceae	ଶର୍ଶକୋଦଙ୍ଗର୍ଜେ (shar-skho-dang-rje)	<i>Heracleum millefolium</i> Diels	QTP-EBT-3202	Food (leaves): seasoning (5)	0.0307
Araceae	ଥ୍ଵୋ (thwo)	<i>Arisaema erubescens</i> (Wall.) Schott	QTP-EBT-4040	1. Food (tubers): mash it and mix it with flour to ferment (12) 2. Animal food (tuber) (2): fodder	0.0920
Araliaceae	ସାନ୍ବର୍ସିଙ୍ଗ (san-bris)	<i>Panax pseudo-ginseng</i> (Burkill)	QTP-EBT-3084	Economic (roots) (29)	0.1779

Table 2 (continued)

Arecaceae	ঝ (spa)	<i>Plectocomia himalayana</i> Griff.	QTP-EBT-4111	Other uses (stems): tools, used to make containers, crutches, etc. (1)	0.0061
Aristolochiaceae	ରିଗେ (ri-gei)	<i>Aristolochia griffithii</i> Hook.f. & Thomson ex Duch.	QTP-EBT-3166	Materials (stems): crafts, the old stems are spiraling and can be used as decorations (1)	0.0061
Asparagaceae	କାଲେମୁଶା (Ka-le-mu-sha), କ୍ରଙ୍କିନ୍ଦ୍ରା (Krung-ki-ndra)	<i>Polygonatum verticillatum</i> (L.) All.	QTP-EBT-3087	1.Medicine (roots): decoction, tonic (2) 2. Economic(roots) (4)	0.0368
Asparagaceae	ଗ୍ୟୋଙ୍କିଲିଯୁଲମ୍ (gyong-khe-lei-bu)	<i>Polygonatum cirrhifolium</i> (Wall.) Royle	QTP-EBT-3156	1.Medicine (roots): decoction, tonic (1) 2. Economic(roots) (1)	0.0123
Balsaminaceae	ଦଙ୍ଗବ୍ୟା (dong-bya)	<i>Impatiens sulcata</i> Wall.	QTP-EBT-3047	1.Food(fruits): fresh fruit (36)2. Animal food (aerial parts) (6)	0.2577
Berberidaceae	ଯୁମୁଶର୍ଶର୍ଶର୍ଶ (Yiu-mu-ser-ser)	<i>Sinopodophyllum hexandrum</i> (Royle)	QTP-EBT-3065	1.Medicine (fruits): raw, gastropathy (6) 2.Food (fruits): fresh fruit (30) 3.Economic (fruits) (17)	0.3252
Berberidaceae	ରିନ୍ବୁ (rin-bu)	<i>Berberis konghoensis</i> Ahrendt	QTP-EBT-3038	1.Medicine(roots): decoction, diarrhea (8) 2.Food (fruits): raw, fresh fruit(8)	0.0982
Betulaceae	ଶରକୁଆ (star-khua)	<i>Betula utilis</i> D.Don	QTP-EBT-3068	1.Medicine(burls):decoction, to regulate blood pressure (6) 2.Other use (stems): making yogurt after peeling (43) 3.Food (stems): poke a hole, drink the water inside (2)4.Fuelwood (sticks) (9)5.Economic (burls) (2)	0.3804
Betulaceae	ଶେଖ୍ବେ (so-skyes), ତେବୀ (tshei-bei)	<i>Corylus ferox</i> Wall.	QTP-EBT-3151	Food (fruits): raw, fresh fruit (5)	0.0307
Boraginaceae	ଅମୁଯୁକୁ (a-mu-yu-klu)	<i>Microula sikkimensis</i> (C.B. Clarke) Hemsl.	QTP-EBT-3034	1.Medicine (leaves): rub with fresh leaves to relieve pain (12) 2.Animal food (whole plants) (3) 3.Food (tender leaves):cooked vegetables (38)	0.3252
Boraginaceae	ଶ୍ରାଙ୍ଗି (smug-tsi)	<i>Onosma hookeri</i> C.B. Clarke	QTP-EBT-3052	1.Medicine (roots): soak it in canola oil, smears, to treat hemorrhoids, acne, inflammation, eczema and promotes hair growth (26) 2.Social uses (roots):the red substance on the roots is embellished on the cakes as a sacrifice for the dead (14)3.Economic (roots) (8)4.Food (flowers):fresh nectar (3)	0.3190
Brassicaceae	ଜାପେଦକା (ja-ped-kha), ପେଦକା (ped-kha), କେଇମୁ (khei-mu)	<i>Thlaspi arvense</i> L.	QTP-EBT-3064	Economic (whole plants) (7)	0.0429

Table 2 (continued)

Pegaeophyton					
Brassicaceae	չՇՌ (chu-ru-skho)	<i>scapiflorum</i> (Hook.f. & Thomson) C.Marquand	QTP-EBT-3016	Food (leaves): cooked vegetable (3)	0.0184
		& Airy Shaw			
Brassicaceae	ՇՌ Շ Ռ (spo-skho-skho), Շ Ռ Ռ Ռ Ռ (spo-spo-lei-htrum)	<i>Capsella bursa-pastoris</i> (L.) Medik.	QTP-EBT-3067	Food(leaves): cooked vegetable(3)	0.0184
Campanulaceae	Ց Ց Ց Ց Ց (blun- bus-kyi-rdo-rje)	<i>Codonopsis foetens</i> Hook.f. & Thomson	QTP-EBT-3029	1.Medicine (roots): decoction, gastropathy (8)2.Economic (roots)(28)3.Food (flowers): fresh nectar (2)	0.2331
Campanulaceae	Չ Չ Չ Չ Չ (chong- klung-me-tog)	<i>Cyananthus lobatus</i> Wall. ex Benth.	QTP-EBT-3058	Medicine (flowers): apply fresh petals to prevent dry lips (28)	0.1718
Campanulaceae	Չ Չ Չ Չ Չ (chong- klung-me-tog)	<i>Cyananthus pedunculatus</i> C.B.Clarke	QTP-EBT-3090	Medicine (flowers): apply fresh petals to prevent dry lips (26)	0.1595
Cannabaceae	Շ Շ Շ Շ Շ (sro-ma-ra-phras)	<i>Cannabis sativa</i> L.	QTP-EBT-3074	Animal food (aerial parts): fodder (5)	0.0307
Caprifoliaceae	Շ Շ Շ Շ Շ (spang-pue)	<i>Nardostachys jatamansi</i> (D.Don) DC.	QTP-EBT-3051	1.Economic (rhizomes) (5) 2.Social uses (rhizomes): ritual use, incense (35)	0.2454
Caprifoliaceae	Շ Շ Շ Շ Շ (spang-tsi- dou)	<i>Pterocephalus hookeri</i> (C.B.Clarke) E.Pritz.	QTP-EBT-3053	Economic (whole plants) (21)	0.1227
Caprifoliaceae	Յ Յ Յ Յ Յ (Yu-mu-thiu-thiu)	<i>Triosteum himalayanum</i> Wall.	QTP-EBT-3083	Food (fruits): raw, fresh fruit (13)	0.0798
Compositae	Շ Շ (go ma)	<i>Taraxacum tibetanum</i> Hand.-Mazz.	QTP-EBT-3020	1.Medicine(whole plants): decoction,used to eliminate fire-evil and treat gynecological diseases (56)2.Food (leaves): cooked vegetable (6) 3.Economic (whole plants) (3)	0.3988
Compositae	Շ Շ Շ Շ Շ (sgang-la-me-tog)	<i>Saussurea tridactyla</i> Sch.Bip. ex Hook.f.	EBT-PL-23	Economic (whole plants) (54)	0.3313
		<i>Leontopodium</i>			
Compositae	Ց (bra)	<i>calocephalum</i> (Franch.) Beauverd	QTP-JPG-0208	Other use (leaves): tool, ignite fire (24)	0.1472
Compositae	Ց Ց Ց Ց Ց (klu-me-tog)	<i>Aster diplostephioides</i> (DC.) C.B.Clarke	QTP-EBT-3056	Economic (whole plants) (23)	0.1472
Compositae	Շ Շ Շ Շ Շ (oh-ma-ji- bu-ji)	<i>Soroseris erysimoides</i> (Hand.-Mazz.) C.Shih	QTP-EBT-3011	1.Veterinary medicine: decoction, eye diseases (1) 2.Animal food (Whole plants)(2) 3.Economic (whole plants) (18)	0.1288
Compositae	Շ Շ Շ Շ Շ (tar-ser-ma)	<i>Cirsium eriophoroides</i> (Hook.f.) Petr.	QTP-EBT-3072	1.Veterinary medicine: decoction, ascaricide (10) 2.Economic (roots) (2) 3.Animal food (whole plants) (7)	0.1166

Table 2 (continued)

Compositae	ဆုန္တိ (ong-ji-ji)	<i>Saussurea topkegolensis</i> H.Ohba & S.Akiyama	QTP-EBT-3021	Economic (whole plants) (15)	0.0982
Compositae	ကြုံမြော (klu-me-tog), ဗျားနိုးနှာ (gyal-po- me-tok)	<i>Aster albescens</i> (DC.) Wall. ex Hand.-Mazz.	QTP-JPG-70834	Economic (whole plants) (12)	0.0736
Compositae	ဟိုမီ (hchi-mei), မြော (klu-me-tog)	<i>Aster flaccidus</i> Bunge	QTP-EBT-3112	Economic (whole plants) (10)	0.0613
Compositae	မြော-ပွဲ (ser-po- khu-hbrus), အော်နှာ (tsh er-ja-me-tok)	<i>Senecio raphanifolius</i> Wall. ex DC.	QTP-EBT-3066	1.Medicine (whole plants): decoction, used to treat cold (3) 2.Economic (whole plants) (6)	0.0552
Compositae	ရှာ (ru-ta), အော်နှာ	<i>Arctium lappa</i> L.	QTP-JPG-102600	1.Medicine (roots): decoction, used to treat cold (5) 2.Other uses (fruits): tools, used to stick mice (1) 3.Animal food (whole plants) (2)	0.0491
Compositae	ဂမာပာ (gams-pa)	<i>Artemisia argyi</i> H.Lév. & Vaniot	QTP-EBT-3154	1.Medicine (aerial parts): bath, used to treat chicken pox (2) 2.Other uses (aerial parts): mosquito repellent (2)	0.0245
Compositae	ခာခုံ (cha-kho-bu)	<i>Cremanthodium reniforme</i> (DC.) Benth.	QTP-EBT-3132	1.Medicine (leaves): smear, used to treat dermatopathy (1) 2. Economic (whole plants) (3)	0.0245
Compositae	ဒ္ဓာ (tsho-sa), ဂမာ	<i>Artemisia younghusbandii</i> J.R.Drumm. ex Pamp.	QTP-EBT-3208	Social uses (aerial parts): incense (3)	0.0184
Compositae	ပြ (bra)	<i>Anaphalis nepalensis</i> (Spreng.) Hand.-Mazz.	QTP-EBT-3123	Economic (whole plants) (1)	0.0061
Compositae	ဂမာလိုင်းရှာ (gams-lin- shar)	<i>Cremanthodium decaisnei</i> C.B.Clarke	QTP-EBT-3132	Economic (flowers) (1)	0.0061
Coriariaceae	ဒ္ဓာ (tso-mo), ဒ္ဓာ	<i>Coriaria terminalis</i>	QTP-EBT-3005	Food (fruits): raw, fresh fruit (7)	0.0429
Crassulaceae	(dong-ru), ဒ္ဓာ	<i>Rhodiola himalensis</i> (D. rje-sro-lo-tmar-po)	QTP-EBT-3017	1.Medicine (roots): slice and decoction, to enhance immunity, regulate blood pressure, treat altitude sickness (8) 2.Economic (roots) (52)	0.3742
Cupressaceae	ရှာ (shug-pa)	<i>Juniperus indica</i> Bertol.	QTP-EBT-3023	Social uses (leaves and stems): ritual use, incense (30)	0.1840
Cupressaceae	ရှာ (spa-ma)	<i>Juniperus tibetica</i> Kom.	QTP-EBT-4000	1.Social uses (stems and leaves): ritual use, incense (5) 2.Food (fruits): raw, fresh fruit (1)	0.0368

Table 2 (continued)

Dennstaedtiaceae	ප්‍රාජමා (pjat-ma)	<i>Pteridium aquilinum</i> (L.) Kuhn	QTP-EBT-3035	Food (tender leaves): cooked vegetables (69)	0.4479
Dryopteridaceae	නුප්‍රාජමා (nu-pjat-ma)	<i>Dryopteris chrysocoma</i> (Christ) C. Chr.	QTP-EBT-3127	Food (tender leaves): cooked vegetable (20)	0.1227
Elaeagnaceae	පිස්කු (pi-skyu)	<i>Elaeagnus umbellata</i> Thunb.	QTP-EBT-3036	Food (fruits): raw, fresh fruit (3)	0.0184
Ephedraceae	ත්සික්ලං (tshi-klung)	<i>Ephedra monosperma</i> J.G.Gmel. ex C.A.Mey.	QTP-JPG-140937	1. Medicine (aerial parts): decoction, used to treat cold (3) 2. Social uses (aerial parts): ritual uses, incense (1) 1. Medicine (roots): soak it in canola oil, use oil to smears hair, cure gray hair (1)	0.0245
Ericaceae	ස්බාං (sba-glung)	<i>Rhododendron anthopogon</i> D. Don	QTP-EBT-4005	2. Social uses (aerial parts): ritual uses, incense(68) 1. Food(leaves): beverage (1) 2. Fuel wood(aerial parts)(35)3. Social uses (aerial parts):ritual uses, incense(4)	0.4294
Ericaceae	සුතු (su-tu)	<i>Rhododendron setosum</i> D. Don	QTP-EBT-3019	1. Food(leaves): beverage (1) 2. Fuel wood(aerial parts)(35)3. Social uses (aerial parts):ritual uses, incense(4)	0.2454
Ericaceae	ඇංජාරු (A-shar-zhar-mo), ඇංජාරු (ji-bu-klung)	<i>Enkianthus deflexus</i> (Griff.) C.K.Schneid.	QTP-EBT-121625	1. Materials (stems): tools, used to stir (3) 2. Food (flowers): raw, nectar (4)	0.0429
Ericaceae	ගේෂුං (gei-shug)	<i>Rhododendron leucaspis</i> Tagg	QTP-IMG-0207	Medicine (flowers): poison (1)	0.0061
Ericaceae	ඡ්‍රොං (zho-shug)	<i>Rhododendron wallichii</i> Hook. f.	QTP-EBT-70828	Fuelwood (aerial parts) (1)	0.0061
Ericaceae	යැංජු (yae-ju)	<i>Rhododendron cinnabarinum</i> Hook. f.	QTP-EBT-4003	Food (flowers): raw (22)	0.1350
Gentianaceae	ස්පංග්-ර්ක්‍යාන්-මේ-තොක් (spang-rkhyan-me-tok)	<i>Gentiana veitchiorum</i> Hemsl.	QTP-EBT-3024	Medicine (whole plants): decoction, used to treat cold, diarrhea, tracheitis (94)	0.5767
Gentianaceae	ඡිංං (ji-zhong)	<i>Gentiana tibetica</i> King ex Hook.f.	QTP-EBT-3057	1. Medicine (leaves): incineration, smear the wound to stop bleeding, veterinary medicine (37) 2. Economic (leaves) (17) 3. Veterinary medicine (leaves) :decoration, epilepsy(4)	0.3558
Gentianaceae	යුලුං-කංජා (yu-lung-khangs-ja)	<i>Halenia elliptica</i> D.Don	QTP-EBT-3059	1. Veterinary medicine (whole plants): decoction, used to treat diarrhea (2) 2. Economic (whole plants) (11)	0.0798
Gentianaceae	තිංහා (tikh-ta), තිංහා (rgya-tikh)	<i>Swertia bifolia</i> Batalin	QTP-EBT-3111	1. Medicine (whole plants): decoction, used to treat cold (5) 2. Veterinary medicine (whole plants): raw, dental ulcer (1) 2. Economic (whole plants) (5)	0.0675

Table 2 (continued)

Geraniaceae	ଲିଗାଦୁ (lig-ga-du)	<i>Geranium nepalense</i> Sweet	QTP-EBT-3061	Economic (whole plants) (2)	0.0123
Grossulariaceae	ରିନ୍-ବୁ (rin-bu)	<i>Ribes glaciale</i> Wall.	QTP-EBT-3082	1.Food(fruit): fresh fruit (36) 2.Animal food (leaves and stems): browse (3)	0.2393
Grossulariaceae	ରିନ୍-ବୁ (rin-bu)	<i>Ribes takare</i> var. <i>desmocarpum</i> (Hook. f. & Thomson) L.T. Lu	QTP-EBT-3037	1.Food (fruit): fresh fruit (25) 2. Animal food (stems and leaves): browse (3)	0.1718
Juglandaceae	ତର୍-କା (tar-kha)	<i>Juglans sigillata</i> Dode	QTP-EBT-3161	Food (Fruits): raw, nut (1)	0.0061
Lamiaceae	ଶର୍କ୍ଷଦଙ୍ଗର୍ଜେ (shar-kh-dang-rje)	<i>Nepeta discolor</i> Royle ex Benth.	QTP-EBT-3135	Food (whole plants): seasoning (33)	0.2147
Lamiaceae	ଲୋ-ମୋ-ମେତୋକ (lo-so-me-tok)	<i>Phlomoides rotata</i> (Benth. ex Hook.f.) Mathiesen	QTP-EBT-3095	1.Medicine (root): incineration, smear to promote wound healing (22) 2.Animal food (aerial parts) (7) 3.Economic (roots) (3) 1.Medicine (whole plants): decoction, gastropathy (11) 2. Food (aerial parts): seasoning (15) 3.Economic (whole plants)(1)	0.1963
Lamiaceae	ଜା (Ja)	<i>Dracocephalum tanguticum</i> Maxim.	QTP-EBT-3207	Medicine (whole plants): decoction,used to treat cold (2)	0.1656
Lamiaceae	ଲୁ-ଲୁ (lu-lu)	<i>Marmoritis complanata</i> (Dunn) A.L.Budantzev	QTP-EBT-3213	Medicine (whole plants): decoction,used to treat cold (2)	0.0123
Lamiaceae	ହା-ଶା-କା (ha-sha-ka)	<i>Nepeta laevigata</i> (D.Don) Hand.-Mazz.	QTP-EBT-3060	Economic (Whole plants) (2)	0.0123
Leguminosae	ଜା-ଶେ-ଖାର (ja-she-khor-khor)	<i>Thermopsis barbata</i> Benth.	QTP-EBT-3101	1.Medicine (stems): decoction, hypertension (1) 2.Animal food (aerial parts): fodder(11) 3.Economic (stems) (2)	0.0859
Liliaceae	ତ୍ସି-କା (tsi-kha)	<i>Fritillaria cirrhosa</i> D.Don	QTP-EBT-3012	1.Medicine (bulbs): boiled in water, used to treat cold and tracheitis (81) 2.Food (fruits): fruit(5) 3.Economic (bulbs) (57)	0.8896
Melanthiaceae	ଚି-ଯୀ-ଯି-ଶ୍ରେ-ହୁଆ (chi-yie-yis-dre-hua)	<i>Paris polyphylla</i> Sm.	QTP-EBT-4060	1.Medicine (whole plants): decoction and smear, used to treat dermatopathy (4) 2.Economic (whole plants) (6) 3.Food(whole plants): cooked vegetable (3)	0.0798
Oleaceae	ତ୍ସି-ପୁ-ଶୁଣ (tsi-pu-shun)	<i>Fraxinus paxiana</i> Lingelsh.	QTP-EBT-3150	1.Medicine (barks): decoction, fracture (49) 2.Veterinary medicine (barks): decoction (4) 3.Economic (barks) (1)	0.3313
Orchidaceae	ବାଙ୍ଗ-ପୋ-ଲକ୍ଷ (bang-po-lakh-)	<i>Gymnadenia conopsea</i> (L.) R.Br.	QTP-EBT-3080	1.Medicine (roots): decoration, cold (18) 2.Economic (roots) (9) 3.Social uses(roots): ritual uses, incense(1)	0.1718

Table 2 (continued)

	pa), Abbreviation:				
	বাংলা (bang-lag)				
Orchidaceae	ঢেহমা (deh-ma)	<i>Gastrodia elata</i> Blume	QTP-JPG-3292	1.Food (rhizomes): used to make soup (2) 2.Economic (rhizomes) (16)	0.1104
Orobanchaceae	ঢেকে ঢেকা (rje-chu-me-tok)	<i>Pedicularis siphonantha</i> D.Don	QTP-EBT-3093	1.Medicine (whole plants): decoction, used to treat cold (3) 2.Economic (whole plants) (2)	0.0307
Papaveraceae	ক্লুমেটো (klu-me-tog)	<i>Meconopsis simplicifolia</i> (D. Don) Walp.	QTP-EBT-3028	Economic (whole plants) (27)	0.1656
Papaveraceae	বেশ্কিয়ুক্লোং (be-skhyu-klong)	<i>Corydalis casimiriana</i> Duthie & Prain ex Prain	QTP-EBT-3054	Economic (root) (1)	0.0061
Pinaceae	শ্বেতক্ষোণ (skhor-khon-la)	<i>Pinus wallichiana</i> A.B.Jacks.	QTB-JL-39	Food (fruits): raw, nut (2)	0.0123
Plantaginaceae	হংলেম (Hung-lem)	<i>Neopicrorhiza scrophulariiflora</i> (Pennell) D.Y.Hong	QTP-EBT-3022	1.Medicine (roots): decoction, used to treat cold (85) 2.Social uses (roots): ritual uses, incense (1) 3.Veterinary medicine (roots): decoction, diarrhea (3) 4.Economic roots) (9)	0.6012
Plantaginaceae	ক্ষেহমা (kheh-ma)	<i>Plantago asiatica</i> L.	QTP-EBT-3117	1.Medicine (whole plants): decoction, used to treat chicken pox (2) 2.Animal food (whole plants): fodder (2) 3.Economic (whole plants) (2)	0.0368
Polygonaceae	চুক্হা (chu-kha)	<i>Rheum nobile</i> Hook. f. & Thomson	QTP-EBT-3010	1.Food (stems): fresh fruits, to be eaten directly after peeling (76) 2.Materials (roots): dye (2) 3.Social uses (leaves): tobacco substitute (1)	0.4847
Polygonaceae	চুশোর (chu-shor)	<i>Rheum acuminatum</i> Hook. f. & Thomson	QTP-EBT-3107	1.Food (stems): fresh fruits, to be eaten directly after peeling (51) 2.Materials (roots): dye:mash it, then boil the water, dye kimchi and clothes (9) 3.Economic (roots) (1)	0.3742
Polygonaceae	পদ্বারালিন (pad-bras-lin-bu)	<i>Polygonum macrophyllum</i> D. Don	QTP-EBT-3116	1.Medicine (roots): decoction, diarrhea(5) 2.Environmental uses (flowers): decoration(1) 3.Animal food (whole plants) (10) 4.Food (seeds): staple food (31) 5. Economic (roots) (5)	0.3129
Polygonaceae	চুক্রি (chu-throe)	<i>Oxyria digyna</i> (L.) Hill	QTP-EBT-3081	1.Medicine (whole plants): decoction, cholecystitis (2) 2.Economic (roots) (1) 3. Food (fruits) (11)	0.0859
Polygonaceae	ন্যালু (nya-lu)	<i>Polygonum tortuosum</i> D. Don	QTP-EBT-3041	1.Medicine (aerial parts): decoction, used to treat diarrhea (2) 2.Animal food (aerial parts): fodder (9)	0.0675

Table 2 (continued)

Polygonaceae	ન્યાલુ (nya-lu), પાંગશારમે- ટોક (pang-shar-me- tok)	<i>Polygonum polystachyum</i> Wall. ex Meisn.	QTP-EBT-3120	1. Food (tender leaves): cooked vegetable (2) 2. Economic (whole plants) (1) 3. Animal food (whole plants): fodder (5) 4. Materials (whole plants): dye (1)	0.0552
Polygonaceae	છુથ્રો (chu-throe), હેલ્બનિ- શોર (he-leh-ni- shor)	<i>Rheum palmatum</i> L.	QTP-JPG-0385	1. Medicine (roots): decoction, used to treat constipate (5) 2. Economic (roots) (3)	0.0491
Primulaceae	ઝોટિક્કાન્ધા (zho-tikh- me-tok)	<i>Primula sikkimensis</i> Hook.	QTP-EBT-3119	1. Environmental use (flowers): season indicators, when flowers bloom, yak milk yields high (7), ornamental plant (10)	0.1043
Primulaceae	હાલોમે- ટોક (ha-lo-me- tok)	<i>Primula concinna</i> Watt	QTP-EBT-3124	Environmental uses (flowers): season indicators, when the flowers bloom, the rainy season arrives (3)	0.0184
Primulaceae	ર્ખાંધાન્ધા (rkhya- khar-me-tok)	<i>Primula denticulata</i> Sm.	QTP-EBT-4047	Environmental uses (flowers): ornamental plant (2)	0.0123
Ranunculaceae	બાંગન્ગા (bang-nga)	<i>Aconitum orochryseum</i> Stapf	QTP-EBT-3165	1. Medicine (roots): decoction, inflammation, diarrhea, cold (44) 2. Veterinary medicine (roots) : decoction, fever and diarrhea (5)	0.2945
Ranunculaceae	ત્સિનથુખ (tsin-thukh)	<i>Aconitum spicatum</i> Stapf	QTP-EBT-3008	1. Medicine (roots): soak in water and apply it to the joints to treat arthritis (17) 2. Veterinary medicine (roots) : decoction, diarrhea (1) 3. Economic (roots) (1)	0.1166
Ranunculaceae	સેરજીમે- ટોક (ser-ji-me- tok)	<i>Caltha palustris</i> L.	QTP-EBT-3043	1. Environmental uses (flowers): when flowers bloom, the quality of ghee is good (1) 2. Economic (whole plants) (2)	0.0184
Ranunculaceae	સુખો (su-khoe)	<i>Thalictrum reniforme</i> Wall.	QTP-EBT-3062	Materials (roots): dye (2)	0.0123
Rosaceae	સેથ્રુ (se-thru)	<i>Rosa omeiensis</i> Rolfe	QTP-EBT-3033	Food (fruit): fresh fruits (65)	0.3988
Rosaceae	અલુંટશેરલું (a-lung- tsher-lung)	<i>Fragaria nubicola</i> (Lindl. ex Hook.f.) Lacaita	QTP-EBT-3049	1. Food (fruits): fresh fruit (49) 2. Economic (stems) (5)	0.3313
Rosaceae	સેલા (Se-la)	<i>Rosa macrophylla</i> var. <i>glandulifera</i> Yu et Ku	QTP-EBT-3003	Food (fruit): fresh fruits (49)	0.2822
Rosaceae	ક્રોમા (kro-ma)	<i>Potentilla anserina</i> L.	QTP-EBT-3055	Food (roots): boiled in water or raw, staple food (45)	0.2638
Rosaceae	સુમુ (su-mu)	<i>Potentilla fruticosa</i> var. <i>arbustula</i> (D.Don) Maxim.	QTP-EBT-3092	Other uses: tool, to make brush (18)	0.1104
Rosaceae	થ્વોંશરલોં (thwo- tsher-long)	<i>Rubus austrotibetanus</i> T.T.Yu & L.T.Lu	QTP-EBT-3006	Food (fruit): raw, fresh fruits (10)	0.0613

Table 2 (continued)

Rosaceae	ཐྲକྲ୍ପୋ (tshar-thkar-po), རྩྭୟ (a-mu-lu)	<i>Sorbus albopilosa</i> T.T.Yu & L.T.Lu	QTP-EBT-3100	1.Tool (roots): used to make farm tools (1) 2. Food (fruits): raw, fresh fruits (2)	0.0184
Rosaceae	ཐྲଶେର୍ପ୍ରମ୍ (tshar-tscher-hprum)	<i>Rosa sericea</i> Wall. ex Lindl.	QTP-EBT-3004	Food(fruits): raw, fresh fruits(3)	0.0184
Rosaceae	ସାଂଶୋ, རାର୍ଦା (cha-shon, Ja-rda)	<i>Sanguisorba diandra</i> (Hook.f.) Nordborg	QTP-EBT-3113	1.Medicine(roots): decoction, used to treat diarrhea (1) 2. Economic (roots) (1)	0.0123
Rutaceae	୩୧୧୧ (thing-ye)	<i>Zanthoxylum oxyphyllum</i> Edgew.	QTP-EBT-4065	Food (Fruits): seasoning (4)	0.0245
Salicaceae	ସେତ (klang-ma)	<i>Salix myrtillacea</i> Andersson	QTP-EBT-3110	Other uses (stems): tools, making yogurt after peeling (19)	0.1166
Salicaceae	ସେତ (klang-ma)	<i>Salix daltoniana</i> Andersson	QTP-EBT-3103	Other uses (stems): tools, making yogurt after peeling (7)	0.0429
Saxifragaceae	ସ୍ମରକ୍ଷଣ (thmar-zhor)	<i>Bergenia purpurascens</i> (Hook.f. & Thomson) Engl.	QTP-EBT-3013	1.Medicine (roots): decoction,diarrhea(37) 2.Veterinary medicine (roots): decoction, diarrhea(13) 3.Economic (roots) (6)	0.3436
Solanaceae	କିନ୍ହିନ୍ଦି (khin-tie), ଜିନ୍ଦି (jin-tin)	<i>Anisodus luridus</i> Link	QTP-EBT-3153	1.Medicine (fruits): smudging, toothache (15) 2. Animal food (whole plants) (2): fodder	0.1043
Tamaricaceae	ମର୍ମ (A-mu)	<i>Myricaria rosea</i> W.W. Sm.	QTP-EBT-3079	Social uses (whole plants): ritual uses, incense, used in funeral rites(2)	0.0123
Taxaceae	ଅଚେନ୍ପା (a-chen-pa)	<i>Taxus wallichiana</i> Zucc.	QTP-EBT-3169	Fuelwood (stems and leaves) (4)	0.0245
Thymelaeaceae	କହରମୁମେଟୋକ (khar-mu-me-tok)	<i>Stellera chamaejasme</i> L.	QTP-EBT-3205	Materials: paper making (roots) (5)	0.0307
Urticaceae	ସାବୁହ୍ରମ (swa-bu-hrum)	<i>Urtica hyperborea</i> Jacq. ex Wedd.	QTP-EBT-3040	1.Food (leaves): cooked vegetables (92) 2.Animal food (aerial parts): fodder (3)	0.5828

plants used to be closely related to the daily life of local people, but now they provide more cultural service.

Discussion

Current situation of the traditional knowledge of plant use
Local Tibetans have a wealth of knowledge of plant utilization, which penetrates into all aspects of life. But these knowledge was mainly distributed among middle-aged people and is severely affected by socio-economic development.

Tibetans have unique dietary habit under special geographical and climatic conditions. Highland barley powder and ghee are the main foods of farmers and herders, and are supplemented by dairy products such as yoghurt and milk residue, as well as meat of mostly beef and lamb [57]. Fruits and vegetables are rare, but in Yadong, many wild vegetables and fruits are used, and these account for

80% of wild edible plants. With the increasing demand for medicinal plants in Tibetan traditional medicine and Chinese medicine, the commercial value of medicinal plants has increased. Yadong County is rich in medicinal plant resources; thus, economic utilisation has become an important category of plant utilisation. In addition, animal husbandry is the main source of livelihood for the local people, and fermented yoghurt plants, animal feed, and seasoning plants related to animal husbandry are also frequently used in Yadong. Herders are the main collectors of high-altitude medicinal plants, which are harvested from alpine meadows and pastures, consistent with ethnobotanical studies in Dolpa, Humla, Jumla, and Mustang districts of Nepal [27].

Experience is an important means for transmitting traditional knowledge [3]. In the past, living conditions were relatively poor, and young people often followed their

Table 3 Informant consensus factor for traditional medicinal plant use categories

Secondary category of use	Tertiary category of use	Number of use reports(Nur)	Number of taxa (Nt)	Informant consensus index factor (FIC)
Cardiovascular disease	Hyperglycaemia, hypertension, anaemia	35	7	0.82
Dermatopathya	Burn, bleeding, acne	116	6	0.96
Gastrointestinal problems	Constipation, diarrhea,gastralgia	118	10	0.92
Infections	Chicken pox	4	2	0.67
Poisons	Poisons	1	1	–
Respiratory complaints	Cold, begma	321	11	0.97
Skeleto-muscular system	Fractures, arthralgia	66	2	0.98
Toothache and mouth	Toothache	15	1	1.00
	Chapped lips	54	2	0.98
Hypoimmunity	Hypoimmunity	2	1	1.00
Inflammations	Inflammations	142	4	0.98
Hair follicle	Promote hair growth, hair darkening	14	2	0.92
Cholecystitis	Cholecystitis	2	1	1.00
Veterinary medicinal	Ulcer, parasites, eyesache, fractures diarrhea	48	10	0.81

elders to gather food, medicine, fuelwood, etc. in the wild. However, the current rapid economic development has resulted in significant supplementation of material resources, and the collection of wild plants has become less necessary. More children are now going to boarding schools in the county and to more developed areas to make a living. Most young people are not interested in traditional medicine, and ethnobotanical studies in some parts of Nepal have described the same phenomenon [28, 58]. Elderly individuals may slowly lose their memory of wild plant use because of socio-economic changes over the years. These reasons have resulted in obstacles to the inheritance of traditional knowledge, and information that has been passed down from generation to generation has become distorted [23, 24]. Ultimately, modern life-style changes are the main reason for the decline in traditional plant knowledge [41].

The most popular edible plants

Wild edible plants (WEPs) play an important role in food supplementation under normal circumstances [58] and are an important source, in addition to cultivated plants, for people to obtain nutrients, vitamins, minerals, and other biologically active compounds [48, 59].

Fruits with high CI values included *Rheum nobile* ($CI_{\text{fruit}} = 0.4663$), *Rosa omeiensis* (0.3988), and *Fragaria*


nubicola (0.3006). The tender stems of *R. nobile* are peeled and eaten raw as snacks by locals. The plant is mainly distributed in the local high mountains at over 4,000 m above sea level [48]. Plants of the same genus are abundant in other Tibetan regions. Tibetans from Lithang eat *R. alexandrae* and *R. palmatum* stems after removing the skin [36]. Tibetans eat the tender stems of *R. officinale* and *R. palmatum* in Zagana, Gansu, China [37]. The Tibetans of Shangri-La, Yunnan, China eat the tender leaves of raw *R. likiangense* [32]. In the Ladakh region, locals eat the young stems of *R. webbianum* [60]. The ripe fruits of *R. omeiensis*, *R. macrophylla* var. *glandulifera* and *R. sericea* are enjoyed by locals, especially children. *R. omeiensis* is also eaten by Tibetans in Sichuan, Gansu, and Shangri-La, Yunnan, China [32, 36, 37]. In addition, the Lhoba people in Douyu village, southeastern Tibet, use it as a medicinal plant to treat anaemia and maintain youth [61]. *F. nubicola* can be eaten raw or made into jams, or eaten with shaved ice. In Nepal Tibetans of the Mustang District also eat this plant as a fresh fruit [36]. Very few gardens and woodlands are available for planting fruit trees in the local area, and the yield of local fruit tree varieties is extremely low. This may be one of the main reasons locals collect wild fruits as a nutritional supplement.

Table 4 Use categories and use reports

The first category	The second category	Criteria	No. of species	Use reports
Food	Fruits	Fruits that were only eaten when they were ripe, such as apple, pear, strawberry	28	534
	Vegetable	Plants material what were used to cook dishes (including making salads directly with raw plant material)	15	408
	Seasoning	Plants that could be added to dishes or soups to increase the flavour of food	7	148
	Starches	Plants that could be used as a direct starch supplement (e.g., tuberous or rhizome of some plants) or processed into starch	3	86
	Beverages	Plants that could be processed into homemade liqueurs or alcoholic beverages and processed into herbal teas	1	1
Economic plant		The living plant, plant part, or derived product that can be traded	53	560
Medicine	Medicine for human	Plants that could be used by local people to treat diseases of human	43	761
	Veterinary medicine	Plants that could be used by local people to treat diseases of animals	10	44
Animal feed	Fodder	Food (herb) for horses and farm animals	16	107
	Browse	Food (leaves of wooden plants) for horses and farm animals	2	6
Social uses	Ritual uses	Plants used in social scenarios, such as incense	11	165
	Somoking substitute	Plants that are substitutes for tobacco	1	1
Fuel Materials	Fuelwood	Wood used for fuel	4	49
	Dyes	Plants that can be used to dye something	3	13
	Crafts	Plants for making crafts, such as wooden bowls	1	1
	Paper making	Raw materials for papermaking	1	1

Table 4 (continued)

The first category	The second category	Criteria	No. of species	Use reports
Environmental uses	Ornamentals	Plants that can be used for ornamental purposes, such as potted plants, head-dresses	3	13
	Season indicators	Plants that can indicate the arrival of the season	3	11
Other uses	Tools	Plants that can be used to make tools, such as containers, cookware	9	117
	Repellent	Plants used to repel mosquitoes	1	2



Urtica hyperborea ($CI_{\text{vegetable}} = 0.5644$) and *Pteridium aquilinum* (0.4294) had vegetables with high CI values. The tender leaves of *U. hyperborea* are locally eaten as wild vegetables. It is consumed in spring and stored as a reserve vegetable during winter. Boiling with rice or tsampa is the main processing method by which locals detoxify the plant. The tender stems of this plant are used to stew soup by Tibetans in Sapi, Ladakh, Jammu and Kashmir, India [36]. A previous study showed that *U. hyperborea* extract lower uric acid levels [62].

Hyperuricaemia and gout affect humans globally [63]. There are abundant resources of *U. hyperborea* in China [64]. Therefore, *U. hyperborea* is expected to be developed into a healthy food source in plateau areas. Young leaves of *P. aquilinum* were collected from the local population; these were blanched and soaked overnight after which they can be cooked as seasonal vegetables. Locals indicated that soaking overnight removes the bitterness and improves the taste of young leaves. However, this plant contains the toxic compound ptaquiloside



Fig. 4 Tibetan cheese fermentation process. **a** *Salix daltoniana* Andersson. **b** *Salix myrtillacea* Andersson. **c** *Betula utilis* D. Don. **d** Soak branches in boiling water, wash, and peel. **e** Cleaned branches. **f** Set up the branches in the bucket, pour the milk, shake the bucket to make the milk adhere to branches. **g** Put the fermented cheese into the pot, add ghee and fry until it is cooked



Fig. 5 The top three medicinal plants. **a** Medicines stored in the home. **b** *Fritillaria cirrhosa* D. Don. **c** *Neopicrorhiza scrophulariiflora* (Pennell) D.Y.Hong. **d** *Gentiana veitchiorum* Hemsl




Fig. 6 The process from collection to utilisation of *Fraxinus paxiana* Lingelsh. **a** the tree. **b** barks. **c** leaves. **d, e** The bark turns the water blue




Fig. 7 Some Tibetan incense plants and incense burner. **a** The incense burner, used to burn Tibetan incense plants. **b** *Juniperus indica* Bertol. **c** *Rhododendron anthopogon* D. Don. **d** *Onosma hookeri* C.B. Clarke. **e** *Juniperus tibetica* Kom. **f** *Nardostachys jatamansi* (D.Don) DC. **g** *Neopicrorhiza scrophulariiflora* (Pennell) D.Y.Hong

(PT), which is carcinogenic [65]. Fortunately, the soaking process removes the carcinogenic toxic substances present in this plant [66]. *P. aquilinum* is also eaten as a wild vegetable by Tibetans in Zhagana, Gansu, China and

Shangri-La, Yunnan, China [32, 37]. In addition, *P. aquilinum* has a high commercial value in Gongba, Gansu, China [35].

The two most popular seasoning species were *Carum carvi* ($CI_{\text{seasoning}} = 0.2270$) and *Nepeta discolor* (0.2147). *C. carvi*, also called “kuo nie”, is the most frequently mentioned spice in Yadong. Local Tibetans collect its young leaves in May or June and fruit in August or September. The fruits of *C. carvi* have a pungent, coriander-like flavour and aroma that originates from essential oils, mostly carvone, limonene, and anethole [67, 68]. *C. carvi* is a source of cumin and caraway seeds, which have been used since ancient times to treat various indications in traditional healing systems in wide geographical areas [69]. Since ancient times, *C. carvi* has been used in Europe as a seasoning spice and an aromatic repellent. It can not only improve eyesight but also make the breath more fragrant. The seeds of this plant are also used as seasoning by Tibetans in Lithang, Sichuan, China and Mustang, Nepal [36]. In addition, the seeds of *C. carvi* is also used a cure for poisoning and fever, promote appetite, and improve digestive health in Lithang, Sichuan, China [70]. The local people collect the aboveground parts of *N. discolor*, and eat them as condiments after drying in the shade.

The locals identified three important starch supplement plants, *Potentilla anserina* ($CI_{\text{starch}} = 0.2638$), *Polygonum macrophyllum* (0.1902). *P. anserina* (Chuoma) is frequently used as a staple in modern Tibet. *P. anserina* roots are boiled and eaten with butter and sugar. This traditional dish is served on important Tibetan holidays [35, 37]. Compared with traditional root foods such as *Solanum tuberosum*, *Ipomoea batatas*, and *Colocasia esculenta*, the ratio of nutrients (protein 9.45%, fat 1.15%, dietary fibre 15.23%) in this plant is healthier and more reasonable [71]. *P. anserina* is also eaten as a substitute for tsampa by the Tibetans Litang in Sichuan, Zhagana in Gansu, and Shangri-La in Yunnan, China [32, 36, 37]. In addition, the local population mix *P. macrophyllum* and *Hordeum vulgare* var. *coeleste* seeds and grind them into flour. Tibetans in Zhagana, Gansu, China use them similarly [37]. Wild starch plants have become an important source of supplementary starches.

These wild plants compensate for the lack of vegetables and fruits in the traditional Tibetan diet and the lack of starch in times of food shortages. In addition, the utilization of these plants also has a high correlation with the surrounding area, which also shows the transmission of culture.

The most popular medicinal plants

We conducted a quantitative analysis based on the information provided by the participants. The five most-cited species were *Gentiana veitchiorum* ($CI_{\text{medicine}} = 0.5767$), *Neopicrorhiza scrophulariiflora* (0.5215), *Fritillaria*

cirrhosa (0.4969), *Taraxacum tibetanum* (0.3436), and *Fraxinus paxiana* (0.3006) (Table 2). As a part of the cultural diversity of Tibetan community, these traditional medicinal knowledge and experiences provided important health services for locals.

It is worth noting that the three most cited medicinal plants are all used to treat respiratory diseases and are usually stocked in the homes of locals to meet daily needs, much like a medical kit for city family (Fig. 5a). Respiratory diseases are the most common diseases locally, which is consistent with ethnopharmacological survey data in Dolpa, Humla, Jumla, and Mustang districts of Nepal [27]. These plant species are all were traditional Tibetan medicines [40, 41] and are used to treat common ailments, such as inflammations, colds, coughs, and diarrhoea. The whole plant body of *G. veitchiorum* (Fig. 5d) is used locally to treat colds and bronchitis, and relevant studies have shown that the plant species have antibacterial, antiviral, and pharmacological activities for treating bronchitis [72–74]. The roots of *N. scrophulariiflora* are widely used by the locals to treat cold (Fig. 5c). According to the Chinese Pharmacopoeia, this plant can treat many diseases [44]. However, there is no documented treatment for the common cold and use as a veterinary medicine. In the Maithili region of eastern Nepal, the plant is used by locals to treat fever and headaches with high consensus [30]. The bulbs of *F. cirrhosa* (Fig. 5b) are used by the locals as a medicine to treat tracheitis, and this species is also rich in pharmacologically active compounds with antitussive activity [75, 76].

T. tibetanum whole plant is added to boiling drinking water, and to treat various inflammations, such as upper respiratory tract infections, and pharyngitis. Young leaves can be collected and used as wild vegetables with a bitter taste. Locals indicated that this vegetable could “clear the heat and remove the fire” and act as a nutritional supplement. Plants of the same genus are also used by Sherpas in Chentang, China, as medicines for cancer and gynaecological diseases, and Tibetans in Shangri-La, China, and Nepal also use it as a wild vegetable [32, 36]. The dandelion plant has a variety of active anti-inflammatory ingredients and contains various nutrients, such as proteins, sugars, vitamins, required by the human body. It is a medicinal and food homologous plant with great developmental value [77].

The bark of *F. paxiana* can be soaked in water to treat fractures and can be used by both humans and livestock. In addition, during the collection process, the locals maintain part of the bark to ensure the nutrient supply of the tree itself, which is also an important manifestation of sustainable collection (Fig. 6). The same usage has been observed in Bhutan [78]. People in Nepal use plants of the same genus to treat body aches [79]. *F. paxiana*

is mainly distributed in the subtropical rainforest below 2,000 m above sea level in the Yadong River Valley, where the population is very small. The main source of *F. paxiana* is Bhutan, where the private sector exchanges and purchases good through trade channels. Local merchants buy the bark of the plant from Bhutan and supply it to drugstores for trading. Although the plant is widely used locally, it has not been documented in traditional Tibetan medicine, and there is no documentation of its efficacy in treating fractures in Chinese medicine [80]. We theorise that the traditional knowledge of this plant was obtained by locals in Bhutan's trade and cultural exchanges.

Economic plant collection and sustainable development

In recent years, Tibetan and Hui medicinal material merchants have collected a large number of medicinal plants in Yadong. The informants informed us that the collection of economic plants had become more intensive compared with approximately 10–20 years ago. However, with the development of commerce, the excessive collection of plants has caused a certain degree of damage to the local ecological environment [28, 34, 81]. The degree of collection and dependence on wild plants is related to the economic status of the local people. It is generally believed that when a certain plant has a high economic value, it may lead to the depletion of plant resources owing to excessive collection [82, 83]. For example, *Fritillaria cirrhosa*, which have high commercial value, has been excessively and indiscriminately harvested (Fig. 3). As a result, its resources are declining sharply, and it is on the verge of extinction [85].

While over-harvesting of some important medicinal plants has increased, many locals are working toward both biological conservation of the medicinal plants through sustainable harvesting and protection of wild species and conservation of their cultural heritage [34]. The local government is also aware of the impact of this uncontrolled gathering on natural vegetation. For example, locals have realized that the harvesting of *Rhodiola himalensis* leads to soil erosion, and the the harvesting of *Rhodiola himalensis* has been banned. *Rhododendron anthopogon*, *Angelica paeoniifolia* and *Fragaria nubicola*, *Rheum palmatum* are cultivated as commercial crops in Kangbu and Shangyadong Township. It is hoped that this will increase local revenues while reducing damage to natural resources.

Although there are many initiatives for sustainable collection and utilisation, these policies have not been fully implemented. As a result, the depletion of high-value medicinal plants that local livelihoods depend on has not been curbed [29].

"Sang" in the daily life of local people

"Sang" is has important social uses and it prevails in Tibetan areas and has a long history [25, 85]. Most socially used plants are commonly used as incense materials, such as *Rhododendron anthopogon* and *Juniperus indica* (Fig. 7), and the dry sticks of these species are burned in a censer, which is placed on the flat roof of a house or at the entrance to the village. Local Tibetans in the Ladakh area also use *J. macropoda* as raw material for Tibetan incense [25]. These plants are usually burned in the early morning to pray for the gods' blessing and good luck for the day.

In ancient society, when the men of the tribe returned after expeditions, hunting, or funerals, people thought that they were contaminated with all kinds of filth. Therefore, their family members used cypress branches and fireworks burned with various herbs to dispel the filth and to prevent its spread, which could, in turn, bring disaster to the family [85]. The following factors influence the plant preferences of local Tibetans. First, most of the plants burned in burner have a fragrance. After these plants are burned, they produce a strong fragrance. The scent drifts in all directions with the wind, so more gods can be reached, thereby fulfilling the needs of gods and providing better protection for the family. Second, most plants used for this ceremony are common people's lives and surroundings.

In Tibetan areas, almost every household is equipped with a mulberry stove (or in the center of the yard, or on the roof by the mountain) (Fig. 7), and this ritual has become an important part of the local peop's live.

Conclusion

This study demonstrates that the diversity of wild plants used by the Tibetan people in Yadong is reflected not only in the number of species but also in the diverse functions of wild plants, including edible plants, medicinal plants, animal feed, social uses, tools, dye, paper making, and other aspects. In this study, reports on the use of edible and medicinal plants were more prominent in the years when modern transportation was underdeveloped, and food supplies were insufficient. Locals have accumulated considerable experience in the use of wild edible plants, which s provide locals with a large amount of nutritional supplements and food supply.

With the development of the social economy, the demand for medicinal materials in the Tibetan and traditional Chinese medicine industries has increased, and the commercial value of many local medicinal plants has been identified. This has brought opportunities for local development, and has a negative impact on the environment. Locals are attempting to use artificial planting methods

to reduce the hazards of overharvesting. Local traditional plant knowledge is also affected by the surrounding areas. This is likely because the Yadong River Valley has been an important trade channel since ancient times, where frequent cultural and trade exchanges have taken place.

In the future, more in-depth research should be conducted on the nutritional components and pharmacological activities of these plants. In addition, resource assessments of local plants with high commercial value should be conducted, and reasonable development strategies should be proposed for species whose survival is significantly threatened.

Acknowledgements

We are very grateful to the informants for sharing their knowledge with us. Extremely gratitude is expressed to MS. Deji for her assistance. We thank Professor Pei Shengji for technical guidance. In addition, we thank Mr. Hai-Kun Xu as auto driver in the wild works, Ms. Hui-Zhao Yang for drawing.

Authors' contributions

WYH organized the study team and provided technical support. GCA and DXY executed the research plan. GCA identified the specimen and wrote the manuscript. GCA, DXY, ZY, ADYW, ZXQ and ZHF collected the data. WYH and DXY reviewed the manuscript. All authors took part in the field works. All authors were involved in the drafting and revision of the manuscript and approved the final revision. All authors read and approved the final manuscript.

Funding

The study was funded by "The Second Tibetan Plateau Scientific Expedition and Research (No. 2019QZKK0502)".

Availability of data and materials

Please contact the corresponding author for data requests.

Declarations

Ethics approval and consent to participate

The authors asked for permission from the local authorities and the people interviewed to carry out the study.

Consent for publication

The people interviewed were informed about the study's objectives and the eventual publication of the information gathered, and they were assured that the informants' identities would remain undisclosed.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Yunnan Key Laboratory for Wild Plant Resources, Kunming Institute of Botany, Chinese Academy of Sciences, 132# Lanhei Road, Yunnan 650201, Kunming, China. ²University of Chinese Academy of Sciences, Beijing, China.

Received: 15 January 2022 Accepted: 7 March 2022

Published online: 31 March 2022

References

1. Maroyi A. Diversity of use and local knowledge of wild and cultivated plants in the Eastern Cape province, South Africa. *J Ethnobiol Ethnomed*. 2017;13(1):43. <https://doi.org/10.1186/s13002-017-0173-8>.
2. Tangjang S, Namsa ND, Aran C, Litin A. An ethnobotanical survey of medicinal plants in the Eastern Himalayan zone of Arunachal Pradesh. *India J Ethnopharmacol*. 2011;134:18–25. <https://doi.org/10.1016/j.jep.2010.11.053>.
3. Eoin LN. Ethnoecology: losing traditional knowledge. *Nat Plants*. 2016;2(8):16125. <https://doi.org/10.1038/nplants.2016.125>.
4. Pei SJ. Review on two decades development of ethnobotany in China. *Plant Divers*. 2008;30(4):505–9.
5. Myers N, Mittermeier RA, Mittermeier CG, Fonseca GABD, Kent J. Biodiversity hotspots for conservation priorities. *NAT*. 2000;403(6772):853–8. <https://doi.org/10.1038/35002501>.
6. Lu XZ. Looking at the prospect of trade development between my country and South Asian countries from the history and current situation of Tibet border trade and Yadong port. *TIB STUD*. 1994;3:8.
7. Zhao YH. *Magical Tibetan culture*. Beijing: THE Ethnic PUBLISHING HOUSE; 2003.
8. DeLancey S: Lhasa Dialect. In *Sino-Tibetan languages* (Vol. 3). Edited by Thurgood G, LaPolla RJ. London: Psychology Press; 2003:270–288.
9. Wang X, Shi SL, Zhang DG. Tibetan grassland nomadic culture (I)—the formation of Tibetans, language and religious beliefs. *Prair lawn*. 2011;31(2):7.
10. Kapstein MT. *The Tibetans*. Oxford: Blackwell Publishing; 2006.
11. Wangyal JT. Ethnobotanical knowledge of local communities of Bumdeling Wildlife Sanctuary, Trashiyangtse. *Bhutan INDIAN J TRADIT KNOW*. 2012;11(3):447–52.
12. Fu Y, Wang Y, Yang YP, Yang XF. Indigenous knowledge and use of *Anisodus tanguticus* in pastoral communities of Eastern Tibet, China, and Its implications for local adaptation. *Plant Divers*. 2015;37(6):881–90. <https://doi.org/10.7677/ynzwyj201515098>.
13. Qiu T, Sun H, Wang HL, Zhou Y, Lou RJ, Yang P, Zhu WT, Sun HB, Du JZ, Zhun G, Jiang SY, Wang XQ. Ethnobotanical study of Juenang cultural area in Rangtang county of northwestern Sichuan, China. *J Chin Mater Med*. 2020;45(3):689–96. <https://doi.org/10.19540/j.cnki.cjcm.20191204.103>.
14. Bhattarai S, Chaudhary RP, Quave CL, Taylor RSL. The use of medicinal plants in the trans-himalayan arid zone of Mustang district. *Nepal J Ethnobiol Ethnomed*. 2010;6:14. <https://doi.org/10.1186/1746-4269-6-14>.
15. Savita LS, Preeti V. Birth related folk medicines of East Sikkim. *Biochem Cell Arch*. 2006;6(2):247–8.
16. Wangchuk P, Pyne SG, Keller PA. An assessment of the Bhutanese traditional medicine for its ethnopharmacology, ethnobotany and ethnoquality: Textual understanding and the current practices. *J Ethnopharmacol*. 2013;148(1):305–10. <https://doi.org/10.1016/j.jep.2013.04.030>.
17. Liu Y, Dao Z, Yang C, Liu Y, Long CL. Medicinal plants used by Tibetans in Shangri-la, Yunnan, China. *J Ethnobiol Ethnomed*. 2009;5(1):15. <https://doi.org/10.1186/1746-4269-5-15>.
18. Kala CP. Indigenous uses, population density, and conservation of threatened medicinal plants in protected areas of the Indian Himalayas. *Conserv Biol*. 2005;19(2):368–78. <https://doi.org/10.2307/3591249>.
19. Salick J, Byg A, Amend A, Gunn B, Law W, Schmidt H. Tibetan medicine plurality. *ECON BOT*. 2006;60(3):227–53. [https://doi.org/10.1663/0013-0001\(2006\)60\[227:TMP\]2.0.CO;2](https://doi.org/10.1663/0013-0001(2006)60[227:TMP]2.0.CO;2).
20. Li Q, Li HJ, Xu T, Du H, ChenLei HG, Fan G, Yi Z. Natural medicines used in the traditional Tibetan medical system for the treatment of liver diseases. *Front Pharmacol*. 2018;9:29. <https://doi.org/10.3389/fphar.2018.00029>.
21. Wang L, Wang YH, Zhuang HF, Zhang Y, Zhang LL, Wang C. The status and characteristics of traditional medicine markets in Dali prefecture. *Plant Divers Resour*. 2013;35(4):8. <https://doi.org/10.7677/ynzwyj201313058>.
22. Fu Y, Wang Y, Yang YP, Yang XF. Indigenous Knowledge and Use of *Anisodus tanguticus* in pastoral communities of Eastern Tibet, China, and Its Implications for Local Adaptation. *Plant Divers Resour*. 2015;37(06):881–90. <https://doi.org/10.7677/ynzwyj201515098>.
23. Li S, Zhang Y, Guo YJ, Yang LX, Wang YH. Monpa, memory, and change: an ethnobotanical study of plant use in Medog County, South-east Tibet, China. *J Ethnobiol Ethnomed*. 2020;16(1):5. <https://doi.org/10.1186/s13002-020-0355-7>.
24. Yang J, Chen WY, Fu Y, Yang T, Luo XD, Wang YH, Wang YH. Medicinal and edible plants used by the Lhoba people in Medog County, Tibet, China. *J Ethnopharmacol*. 2020;249: 112430. <https://doi.org/10.1016/j.jep.2019.112430>.
25. Bhattacharyya A. Ethnobotanical observations in the Ladakh region of northern Jammu-and-Kashmir State, India. *Econ Bot*. 1991;45(3):305–8. <https://doi.org/10.1007/BF02887067>.

26. Bhattacharai S, Chaudhary RP, Taylor RSL. Ethnomedicinal plants used by the people of Manang district, central Nepal. *J Ethnobiol Ethnomed*. 2006;2:41. <https://doi.org/10.1186/1746-4269-2-41>.

27. Kunwar RM, Nepal BK, Kshhetri HB, Rai SK, Bussmann RW. Ethnomedicine in Himalaya: a case study from Dolpa, Humla, Jumla and Mustang districts of Nepal. *J Ethnobiol Ethnomed*. 2006;2:27. <https://doi.org/10.1186/1746-4269-2-27>.

28. Singh AG, Kumar A, Tewari DD. An ethnobotanical survey of medicinal plants used in Terai forest of western Nepal. *J Ethnobiol Ethnomed*. 2012;8:19. <https://doi.org/10.1186/1746-4269-8-19>.

29. Kunwar RM, Subedi BP, Baral SR, Maraseni T, LeBoa C, Adhikari YP, Bussmann RW. Ethnobotany of the Himalayas: the Nepal, Bhutanese, and Tibetan Himalayas. *Ethnobot Mt*. 2020;09:1–39.

30. Kunwar RM, Fadiman M, Cameron M, Bussmann RW, Thapa-Magar KB, Rimal B, Sapkota P. Cross-cultural comparison of plant use knowledge in Baitadi and Darchula districts. *Nepal Himalaya J Ethnobiol Ethnomed*. 2018;14:40. <https://doi.org/10.1186/s13002-018-0242-7>.

31. He J, Yang B, Dong M, Wang Y. Crossing the roof of the world: trade in medicinal plants from Nepal to China. *J Ethnopharmacol*. 2018;224:100–10. <https://doi.org/10.1016/j.jep.2018.04.034>.

32. Ju Y, Zhuo JX, Liu B, Long CL. Eating from the wild: diversity of wild edible plants used by Tibetans in Shangri-la region, Yunnan, China. *J Ethnobiol Ethnomed*. 2013;9(1):28. <https://doi.org/10.1186/1746-4269-9-28>.

33. Ding XY, Zhang Y, Wang L, Zhuang HF, Chen WY, Wang YH. Collection calendar: the diversity and local knowledge of wild edible plants used by Chenthang Sherpa people to treat seasonal food shortages in Tibet. *China J Ethnobiol Ethnomed*. 2021;17(1):40–40. <https://doi.org/10.1186/s13002-021-00464-x>.

34. Bhattacharai S, Chaudhary RP, Quave CL, Taylor RSL. The use of medicinal plants in the trans-himalayan arid zone of Mustang district. *Nepal*. 2010;6:14. <https://doi.org/10.1186/1746-4269-6-14>.

35. Kang YX, Łuczaj Ł, Kang J, Wang F, Hou JJ, Guo QP. Wild food plants used by the Tibetans of Gongba Valley (Zhouqu county, Gansu, China). *J Ethnobiol Ethnomed*. 2014;10:20. <https://doi.org/10.1186/1746-4269-10-20>.

36. Boesi A. Traditional knowledge of wild food plants in a few Tibetan communities. *J Ethnobiol Ethnomed*. 2014;10(1):75. <https://doi.org/10.1186/1746-4269-10-75>.

37. Kang J, Kang YX, Ji XL, Guo QP, Guillaume J, Marcin P, Nasim Ł, Li DW, Łukasz Ł. Wild food plants and fungi used in the mycophilous Tibetan community of Zhagana (Tewo County, Gansu, China). *J Ethnobiol Ethnomed*. 2016; 12: 21. <https://doi.org/10.1186/s13002-016-0094-y>.

38. Zhang J, Longzhu DJ, Kang JH, Sheng Z, La B. Ethnobotanical study on Tibetan substituting tea plants in Banma Area. *Qinghai Chin Wild Plant Resour*. 2020;39(8):80–5.

39. Savita LS, Preeti V, Parveen S. Studies on wild edible plants of ethnic people in east Sikkim. *Asian J BioSci*. 2006;1(2):117–25.

40. Li FF, Zhuo JX, Liu B, Devra J, Long CL. Ethnobotanical study on wild plants used by Lhoba people in Milin County. *Tibet J Ethnobiol Ethnomed*. 2015;11(1):21. <https://doi.org/10.1186/s13002-015-0009-3>.

41. Thakur D, Sharma A, Uniyal SK. Why they eat, what they eat: patterns of wild edible plants consumption in a tribal area of Western Himalaya. *J Ethnobiol Ethnomed*. 2017;13:70. <https://doi.org/10.1186/s13002-017-0198-z>.

42. Yangjin ZG, Tu YL, Wen XM. An ethnobotanical study on the utilization of cedar plants by Tibetans. *Tibet Technol*. 2013;000(012):71–3. <https://doi.org/10.3969/j.issn.1004-3403.2013.12.031>.

43. Savita LS, Preeti V. Plants used as tooth-brushes by ethnic people of East Sikkim. *Adv Plant Sci*. 2006;19(2):561–2.

44. La Z. Analysis of tourism meteorological conditions in yadong county. *Agr Technol Serv*. 2017;34(016):58–58. <https://doi.org/10.3969/j.issn.1004-8421.2017.16.049>.

45. Ji XP. Practice and thinking of urbanization construction in Yadong County. *Tibet Theoretic Platf Tibetan Devel*. 2013;01(1):58.

46. Can MY, Jiu M, Deji CM, Zhang XQ, Bao YH. Investigation on development status and business model of Pali yak industry in Yadong County. *Tibet Agric Sci*. 2020;42(4):4.

47. Yu KF. On the county construction and port economy in Yadong. *Tibet World Regional Stu*. 1998;1:86–9.

48. Wu Z, Raven PH, Hong D. *Flora of China*. Beijing: Science Press. St. Louis: Missouri Botanical Garden Press; 1994.

49. The Plant List. Version 1.1. Published on the internet. <http://www.theplantlist.org/>, Accessed 1st Jan 2013.

50. Reyes-García V, Huanca T, Vadez V, Leonard W, Wilkie D. Cultural, practical, and economic value of wild plants: A quantitative study in the Bolivian Amazon. *Econ Bot*. 2006;60(1):62–74. [https://doi.org/10.1663/0013-0001\(2006\)60\[62:cpevo\]2.0.co;2](https://doi.org/10.1663/0013-0001(2006)60[62:cpevo]2.0.co;2).

51. Cook F. *International Databases for Plant Sciences (TDWG)*. Economic botany data collection standard. Kew: Royal Botanic Gardens; 1995.

52. Tardío J, Pardo-de-Santayana M. Cultural importance indices, a comparative analysis based on the useful wild plants of Southern Cantabria Northern Spain. *Econ Bot*. 2008;62:24–39. <https://doi.org/10.1007/s12231-007-9004-5>.

53. Troter R, Logan M. Informant consensus: a new approach for identifying potentially effective medicinal plants. In: Etkin NL, editor. *Indigenous medicine and diet: biobehavioural approaches*. New York: Redgrave Bedford Hills; 1986. p. 91–112.

54. Zang X, Gamacuo, Zhadangzhou. A sociological study of religion on the gender division of labor among Tibetan herdsman. *Guizhou Ethnic Stud*. 2013; <https://doi.org/10.13965/j.cnki.gzmzyj10026959.2013.01.024>.

55. Alexis DR, Ana IM, Alejandro C, Alicia C, Andrés C. Traditional climate knowledge: a case study in a peasant community of Tlaxcala. *Mexico J Ethnobiol Ethnomed*. 2016;12:33. <https://doi.org/10.1186/s13002-016-0105-z>.

56. Kang J, Kang Y, Feng J, Liu M, Ji X, Li D, Stawarczyk K, Łuczaj Ł. *J Ethnobiol Ethnomed*. 2017;13(1):1–14. <https://doi.org/10.1186/s13002-017-0165-8>.

57. Jia L. On the changes of tibetan food culture customs. *J Manag*. 2009;5X:1.

58. Mallik AR, Chaudhary S, Shrestha S. Useful valuable plants of Maithili community in eastern Nepal: an ethnobotanical study. *Bangl J Plant Taxon*. 2020;27(2):439–46. <https://doi.org/10.3329/bjpt.v27i2.50678>.

59. Ogle BM, Hung PH, Tuyet HT. Significance of wild vegetables in micronutrient intakes of women in Vietnam: an analysis of food variety. *Pac J Clin Nutr*. 2001;10(1):21–30. <https://doi.org/10.1046/j.1440-6047.2001.00206.x>.

60. Ballabh B, Chaurasia O, Pande P, Ahmed Z. Raw edible plants of cold desert Ladakh. *Indian J Tradit Know*. 2007;6(1):182–4. <https://doi.org/10.1007/s11627-006-9012-7>.

61. Chen WY, Yang T, Yang J, Qiu ZC, Ding XY, Wang YH, Wang YH. Wild plants used by the Lhoba people in Douyu Village, characterized by high mountains and valleys, in southeastern Tibet, China. *J Ethnobiol Ethnomed*. 2021;17(1):46. <https://doi.org/10.1186/s13002-021-00472-x>.

62. Han S, Wei R, Han D, Zhu J, Zhong G. Hypouricemic effects of extracts from *urtica hyperborea* Jacq. ex Wedd. in hyperuricemia mice through XOD, URAT1, and OAT1. *Biomed Res Int*. 2020;11:1–8. <https://doi.org/10.1155/2020/2968135>.

63. Yang H, Gao L, Niu Y, Zhou Y, Lin H, Jiang J, Kong X, Liu X, Li L. Mangiferin inhibits renal urate reabsorption by modulating urate transporters in experimental hyperuricemia. *Biol Pharm Bull*. 2015;38(10):1591–8. <https://doi.org/10.1248/bpb.b15-00402>.

64. Chen X, He S, Lu Y, Yuan L, Zhang Q. Inhibition of spontaneous canine benign prostatic hyperplasia by an *urtica fissa* polysaccharide fraction. *Planta Med*. 2014;81(1):10–4. <https://doi.org/10.1055/s-0034-1383364>.

65. Rasmussen L, Lauren D, Smith B, Hansen H. Variation in ptaquiloside content in bracken (*Pteridium esculentum* (Forst f) Cockayne) in New Zealand. *New Zeal Vet J*. 2008;56(6):304–9. <https://doi.org/10.1080/00480169.2008.36851>.

66. Liu Y, Wujiguleng W, Long CL. Food uses of ferns in China: a review. *Acta Soc Bot Pol*. 2012;81(4):263–70. <https://doi.org/10.5586/asbp.2012.046>.

67. López MD, Jordán MJ, Pascual-Villalobos MJ. Toxic compounds in essential oils of coriander, caraway and basil active against stored rice pests. *J Stored Prod Res*. 2008;44(3):273–8. <https://doi.org/10.1016/j.jspr.2008.02.005>.

68. Baanou S, Bagdonaitė E, Marongiu B, Piras A, Porcedda S, Falconieri D. Extraction of the volatile oil from *Carum carvi* of Tunisia and Lithuania by supercritical carbon dioxide: chemical composition and antiulcerogenic activity. *Nat Prod Res*. 2013;27(22):2132–6. <https://doi.org/10.1080/1476419.2013.771350>.

69. Deepak. Importance of *Cuminum cyminum* L and *Carum carvi* L in traditional medicaments - a review. *Indian J Tradit Know*. 2013; 12(2): 300–307. [https://doi.org/10.1016/S0009-2614\(03\)01204-1](https://doi.org/10.1016/S0009-2614(03)01204-1).

70. Sun ML, Jiang ZT, Li R. Research progress and application of natural seasoning spice *Carum carvi* L. *Chin Condiment*. 2009;34(365):24–6. <https://doi.org/10.3969/j.issn.1000-9973.2009.07.002>.

71. Sun J, Lv JP, Bo HB. Analysis and evaluation of nutrient components of Tibetan medicine *Potentilla anserina* L. *Food Sci.* 2008;29(2):4. <https://doi.org/10.3321/j.issn.1002-6630.2008.02.092>.
72. Liu JJ, Li XB, Geng Z, Zhang X, Hou Y, Liu SB, Xu WZ, Tian Q. Test of *Gentiana veitchiorum* Hemel against MRSA. *Chin Pharmacol Bull.* 2011;27(7):1024–7. <https://doi.org/10.3969/j.issn.1001-1978.2011.07.031>.
73. Wei PF, Tian Q, Li XB, Guo WP, Li XB, Liu JJ, Chen JJ. Preliminary experimental study on anti-respiratory syncytial virus of *Gentiana veitchiorum* Hemel extract. *J Pediatr Pharm.* 2011;17(2):4.
74. Geng Z, Li XB, Hou Y, Liu JJ, Liu SB, Tian Q. Study on the effective components of *Gentiana veitchiorum* Hemel in the treatment of chronic bronchitis in mice. *J Chin Med Mater.* 2011;17(2):4. <https://doi.org/10.13863/j.issn1001-4454.2010.03.041>.
75. Zhou Y, Ji H, Li P, Jiang Y. Antimuscarinic function of five fritillaria alkaloids on Guinea Pig tracheal tracheal strips. *J China Pharm Univ.* 2003;34(1):58–60. <https://doi.org/10.3321/j.issn.1000-5048.2003.01.015>.
76. Li P, Ji H, Xu GJ, Xu LS. Antitussive and expectorant effect of Fritillaria traditional Chinese medicine. *J China Pharm Univ.* 1993;24(6):3.
77. Schutz K, Carle R, Schieber A. Taraxacum - A review on its phytochemical and pharmacological profile. *J Ethnopharmacol.* 2006;107(3):313–23. <https://doi.org/10.1016/j.jep.2006.07.021>.
78. Wangchuk P, Yeshi K, Jamphey K. Pharmacological, ethnopharmacological, and botanical evaluation of subtropical medicinal plants of Lower Kheng region in Bhutan. *Integr Med Res.* 2017;6(4):372–87. <https://doi.org/10.1016/j.imr.2017.08.002>.
79. Joshi K. Indigenous knowledge and uses of medicinal plants in Macchegaun Nepal. *Indian J Tradit Know.* 2011;10(2):281–6. <https://doi.org/10.1139/B11-014>.
80. Sun SS. On the original plant of Chinese herbal medicine "Qinpí". *Bull.Med-Ethno-Bot.Res.* 1998; 01: 51–88. doi: CNKI: SUN: MBZW.0.1988-01-003.
81. Zhang XZ, Yang YP, Piao SL, Bao WK, Wang GX. Ecological change on the tibetan plateau. *Chin J.* 2015;60(32):3048. <https://doi.org/10.1360/N972014-01339>.
82. Primack R, Ji WZ. A primer of conservation biology. Beijing: China Forestry Publishing House; 2000.
83. Williams VL, Balkwill K, Witkowski ETF. Unraveling the commercial market for medicinal plants and plant parts on the Witwatersrand. *South Africa Econ Bot.* 2000;54:310–27. <https://doi.org/10.1007/BF02864784>.
85. Guo SL. A review of the research on the cherished medicinal plant *Fritillaria cirrhosa*. *Tibet Technol.* 2020;12:2. <https://doi.org/10.3969/j.issn.1004-3403.2020.12.006>.
85. Li ML, Xu JC. The "Wei sang" custom of Tibetan families in Yunnan——Taking two Tibetan communities in Diqing Tibetan Autonomous Prefecture as an example. *Ethno- nat'l. Studies.* 2007; 169 (06): 46–55.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

